

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Serial No. : 09/836,705  
Applicants : Yuki ABE et al.  
Filed : April 17, 2001  
Art Unit : 1652  
Examiner : Kathleen M. Kerr  
Docket No. : 01149/HG  
Customer No. : 01933  
Confirmation No. : 7090

Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

DECLARATION UNDER 37 CFR 1.132

S I R :

I, Yuki ABE, declare as follows:

1. I am a coinventor of the above-identified patent application.

2. I graduated from the University of Tokyo, Japan in 1996, and I received a Master's degree in the field of agriculture.

3. I have worked for Sankyo Company, Limited of Tokyo, Japan, since 1996, and I presently a researcher.

4. The following information was generated by experiments that were carried out under my supervision.

Two panels are attached hereto. The two panels show the results of genomic southern blot hybridizations with the *mlcR* and the *mlcE* cDNA as probes, respectively. The hybridizations were performed according to Example 5 on pages 42 to 44 of the specification of application Serial No. 09/836,705, the entire contents of which are hereby incorporated by reference herein.

The genomic DNA of *Penicillium citrinum* SANK 13380, *Penicillium chrysogenum*, *Penicillium notatum*, *Penicillium italicum* and *Eupenicillium sinaicum* were digested separately with a restriction enzyme BamH 1 at 37°C for 2 hours and then loaded on lanes 1 to 5 of each gel, respectively.

Only one band was specifically detected in each lane of the left panel, and two bands were specifically detected in each lane of the right panel.

All of the *Penicillium* species *chrysogenum*, *notatum* and *italicum* and the *Eupenicillium* species *sinaicum* were confirmed to produce ML-236B and can be distinguished from *Penicillium citrinum* on the basis of the morphological and taxonomical analysis.

These results demonstrate that there are genes corresponding to *mlcR* and *mlcE* in the genome of ML-236B producing microorganisms and said genes contribute to the production of ML-236B in said microorganisms, and further that ML-236B-producing microorganisms other than *Penicillium citrinum* can be used as a host cell in the claimed methods.


I hereby declare that all statements made herein of my own knowledge are true, and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or

imprisonment, or both, under Section 1001, of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.


Date: October 6, 2003

By:


Yuki Abe  
Yuki ABE



*mlcR*



*mlcE*



1. *Penicillium citirinum*
2. *Penicillium chrysogenum*
3. *Penicillium notatum*
4. *Penicillium italicum*
5. *Eupenicillium sinaicum*

143

V-54

# MYCOTAXON

AN INTERNATIONAL JOURNAL DESIGNED TO EXPEDITE PUBLICATION  
OF RESEARCH ON TAXONOMY & NOMENCLATURE OF FUNGI & LICHENS

---

A NEW EUPENICILLIUM SPECIES WITH  
RETICULATELY ORNAMENTED ASCOSPORES

SHUN-ICHI UDAGAWA AND SEIICHI UEDA

Reprinted from MYCOTAXON 14: 266-272. 1982.

Published 22 January 1982

# MYCOTAXON

Vol. XIV, No. 1, pp. 266-272

January-March 1982

## A NEW EUPENICILLIUM SPECIES WITH RETICULATELY ORNAMENTED ASCOSPORES

Shun-ichi UDAGAWA

*National Institute of Hygienic Sciences, Kamiyoga 1-  
chome, Setagaya-ku, Tokyo 158, Japan*

AND

Seiichi UEDA

*Nagasaki Prefectural Institute of Public Health and  
Environmental Sciences, Nameshi 1-chome,  
Nagasaki 852, Japan*

Members of *Eupenicillium*, one of several groups of ascomycetes having *Penicillium* anamorphs, have been reported to be of wide-spread occurrence throughout the world (Raper and Thom, 1949; Scott, 1968; Pitt, 1979). As with *Talaromyces* species, field soils apparently provide *Eupenicillium* spp. with their most common habitats. They are also frequently encountered in marine environments. Ueda (1980, and a paper presented at the Annual Meeting of the Mycological Society of Japan, held at Niigata, May 28th, 1981) reported a number of *Eupenicillium* species from marine sludges in Nagasaki Bay, southern Japan, and suggested that their presence there might be attributed to sedimentation from surrounding terrestrial habitats. *Eupenicillium brefeldianum* (B.O. Dodge) Stolk & Scott and *E. javanicum* (van Beyma) Stolk & Scott are the most commonly encountered species from marine and estuarine environments.

A member of the genus with reticulately ornamented ascospores was isolated during a continuing study of ascomycetous microfungi in marine sludges. It proved to be sufficiently different from all described taxa of the genus to warrant its description as a new species.

*Eupenicillium sinaicum* Udagawa & Ueda, sp.nov. (Figs. 1-4,  
6)  
Stat. anam. *Penicillium sinaicum* Udagawa & Ueda, st.  
nov.





Coloniae in agaro Czapekii satis crescentes, sulcatae vel plus minusve rugosae, e coacto crasso mycelio compositae, floccosae, albae vel roseo-bulbalinae vel roseo-vinosae, numerosis obscuris cleistotheciis et sparsis conidiilibus fructificationibus formantes; reversum roseo-bulbalinum vel incarnatum. Coloniae in agaro maltoso effusae, tenues, floccosae vel velutinae, deinde granulosaе, croceae; cleistothecia abunde producentia et conidia sparsa; reversum melleum vel isabellinum. Coloniae in agaro farina avenae mixto effusae, tenuiores, granulosaе, salmoneae; cleistothecia abunde producentia; reversum incolor vel aliquantum ferrugineum.

Cleistothecia superficialia, dispersa vel confluentia, dilute flavo-brunnea, subglobosa vel elongata, 150-280  $\mu\text{m}$  diam, glabra vel hyphis pigmentiferis laxè intricatis circumdata; peridium primo sclerotioideum, deinde tenue et membranaceum. Asci catenati in ascogonicis hyphis, octospori, subglobosi vel ovati, 9-11.5  $\times$  8-10  $\mu\text{m}$ , postremo evanescentes. Ascosporae hyalinae vel dilute flavae, late lenticulares, 3.0-4.0  $\times$  2.5-3.0  $\mu\text{m}$ , cum subtili aequatoriali sulco praeditae sed sine distincte cristis, reticulatis convexis ornatae (sub 'SEM').

Penicilli monoverticillati vel plerumque biverticillati et asymmetrîci. Conidiophora hyalina, levîa, septata, variabilia, e substrato erecto 350-450(-900)  $\times$  2.5-3.0 (-3.5)  $\mu\text{m}$ , vel e hyphis aeriis oriunda usque 80  $\mu\text{m}$  longa. Rami interdum praediti, 25-35  $\mu\text{m}$  longi. Metulae 2-4 per verticillum, 12.5-15(-20)  $\times$  2.5-4.0  $\mu\text{m}$ , hyalinae, leves. Phialides 3-4 per verticillum, 9-12.5  $\times$  2.0-2.5  $\mu\text{m}$ . Conidia hyalina, globosa vel subglobosa, 2.0-3.0  $\mu\text{m}$  diam, levîa, catenata. Chlamydosporae abunde producentes, vulgo intercalares.

Holotypus No. 2894, NHL, isolatus e limo mari in 'Port Said', Aegypto, 9.x.1978, leg. H. Komatsu. In herb. NHL.

Etymology: lat. *sinaicum*, referring to the Sinai peninsula, Egypt, the type locality.

Colonies on Czapek agar growing moderately, attaining a diameter of 4.0 cm in 14 days at 23 C, radially furrowed, more or less wrinkled, consisting of a thick mycelial felt with floccose surface, in white to pale pink colors near Rosy Buff (Rayner, 1970) or Rosy Vinaceous (Rayner, 1970), developing a dense layer of cleistothecia on the felt, obscured by a loose overgrowth of aerial hyphae; conidia-bearing structures limited and not affecting the colony appearance; exudate lacking; odor slightly musty; reverse dull orange (Rosy Buff or Flesh; Rayner, 1970) with a deep red tint.



Colonies on malt extract agar growing rapidly, attaining a diameter of 4.2-4.6 cm in 14 days at 23 C, thin, more or less furrowed, floccose to velvety, with surface becoming granular by the development of abundant cleistothecia, in pale brown colors near Saffron (Rayner, 1970), with limited production of conidia; exduate lacking; odor indistinct; reverse pale yellowish brown near Honey or Isabeline (Rayner, 1970).

Colonies on oatmeal agar growing as on malt but thinner, granular by the development of abundant cleistothecia, with limited conidia, in pale yellow-orange colors near Salmon (Rayner, 1970); exduate abundant at submarginal areas, clear; odor lacking; reverse uncolored or somewhat dull reddish orange (Rust; Rayner, 1970).

Cleistothecia superficial, scattered or aggregated in small clusters, pale yellowish brown, subglobose to elongate, 150-280  $\mu\text{m}$  in diam, glabrous or loosely covered with pigmented hyphae, maturing rather slowly from the center outwards, within 21 days. Peridium at first sclerotoid, consisting of masses of thick-walled, polygonal cells, later becoming thin and membranaceous. Asci typically borne in curved chains which consist of up to 6-7 cells, arising as branches from ascogenous hyphae, 8-spored, subglobose to ovate, 9-11.5  $\times$  8-10  $\mu\text{m}$ , evanescent at maturity. Ascospores hyaline to pale yellow, broadly lenticular, 3.0-4.0  $\times$  2.5-3.0  $\mu\text{m}$ , with equatorial areas flattened to show a shallow furrow but distinct crests lacking, with convex surfaces distinctly reticulate under SEM observation.

Penicilli (conidia-bearing structures) monoverticillate to mostly biverticillately asymmetrical. Conidiophores hyaline, smooth-walled, septate, variable, arising either from the substratum, 350-450(-900)  $\times$  2.5-3.0(-3.5)  $\mu\text{m}$ , or as short branches from aerial hyphae mostly less than 80  $\mu\text{m}$  in length. Branches sometimes present, 25-35  $\mu\text{m}$  in length. Metulae commonly in groups of 2-4, measuring 12.5-15(-20)  $\times$  2.5-4.0  $\mu\text{m}$ , hyaline, smooth-walled. Phialides flask-shaped, 3-4 in the verticil, 9-12.5  $\times$  2.0-2.5  $\mu\text{m}$ . Conidia hyaline, globose to subglobose, 2.0-3.0  $\mu\text{m}$  in diam, smooth-walled, connective in tangled or loosely column chains. Chlamydospores abundantly produced, mostly intercalary.

At 37 C, growth is nil.

Isolation: marine sludge, the Suez Canal, 30 km north from Port Said, Sinai Peninsula, Egypt, 9.x.1978, coll. H. Komatsu, No. 2894, NHL (holotype).

In his monographic studies, Scott (1968) pointed out that, on the basis of ascus arrangement within the cleisto-





Fig. 1. *Eupenicillium sinaicum*, NHL 2894.  
 A. Asci. B. Ascospores. C. Penicilli. D. Conidia.

thecia, the species of *Eupenicillium* can be divided into two distinct series: one with species producing catenate asci, and other with asci borne singly. A scanning electron microscopic observation of the ascospore ornamentation for 21 *Eupenicillium* species was given by Udagawa and Horie (1973a), who proposed that the ascospore morphology should be the primary criterion for species separation. They separated the genus into two major groups of species, those bearing distinct equatorial crests and those without such ornamentation.

Pitt (1979), however, stated that 'such characters sometimes appear to have little relationship with other characters considered to be important in the classification of similar fungi, such as growth rates, colony morphology and anamorphic states'. We support the opinion that morphologies for all reproductive states should be consulted at the species level. Nevertheless, in the taxonomy of *Eupenicillium*, the above features of asci and ascospores provide a useful basis for species separation. As an easily discernible and very stable feature, ascospore morphology traditionally has been used in the identification of species in other genera with cleistothecial ascocarps, particularly in the Eurotiaceae, e.g. *Eurotium*, *Neosartorya*, *Talaromyces*, etc. In our experience, growth rates of *Eupenicillium* species, even under standard conditions, are often unreliable. Conidia-bearing structures are sometimes lacking in fresh isolates of *Eupenicillium* species and are extremely inconspicuous or fragmentary in some strains of *E. ornatum* Udagawa, *E. meloforme* Udagawa & Horie, *E. parvum* (Raper & Fennell) Stolk & Scott, *E. rubidum* Udagawa & Horie, etc. (Udagawa, 1968; Udagawa and Horie, 1973b).

*Eupenicillium sinaicum* becomes the fourth species of the genus known to produce ascospores with a reticulate wall. *Eupenicillium molle* Malloch & Cain (1972) has asci disposed in curved or helicoid chains, and appears to be the most closely related species. The ascospores of both of these species have nearly the same range in length and width. However, *E. molle* is clearly different from this species: the asci are smaller; the ascospores have 2 low equatorial ridges and a more irregular ornamentation (as can be seen by comparing Figs. 2, 3, 5, and Fig. 8 in the paper of Udagawa and Horie, 1973a); the penicilli vary in complexity from biverticillate, composed of terminal verticils of phialides numbering up to 8 or 10, to more variously branched and terverticillate, but less commonly have the monoverticillate pattern of *E. sinaicum*.





Figs. 2-4 and 6. *Eupenicillium sinaicum*, NHL 2894.  
2, 3. SEM micrographs of ascospores. 4. Asci. 6. Penicilli. Fig. 5. SEM micrograph of ascospores of *Eupenicillium molle* Malloch & Cain, IMI 84589.

(scale-lines: 2=5  $\mu$ m, 3=2  $\mu$ m, 4=10  $\mu$ m, 5=3  $\mu$ m and 6=50  $\mu$ m.)

The reticulate ascospore ornamentation of *E. sinaicum* is also somewhat similar to that of *E. meloforme* Udagawa & Horie (1973b) and *E. reticulisporum* Udagawa (1968) but differs in the lack of equatorial ridges. This species is primarily distinguished from the latter two species by the formation of its asci in helicoid chains.

#### ACKNOWLEDGMENT

The authors would like to thank Professor D. Malloch of the University of Toronto for critically reviewing the manuscript.

#### REFERENCES

- Malloch, D., and R. F. Cain. 1972. New species and combinations of cleistothecial Ascomycetes. *Can. J. Bot.* 50: 61-72.
- Pitt, J. I. 1979. The genus *Penicillium* and its teleomorphic states *Eupenicillium* and *Talaromyces*. Academic Press, London. 634 p.
- Raper, K. B., and C. Thom. 1949. A manual of the Penicillia. Williams & Wilkins Co., Baltimore. 875 p.
- Rayner, R. W. 1970. A mycological colour chart. Commonwealth Mycological Institute, Kew.
- Scott, D. B. 1968. The genus *Eupenicillium* Ludwig. Microbiol. Res. Group, Bull. 1, Council for Scientific and Industrial Research, South Africa. Pretoria. 150 p.
- Udagawa, S. 1968. Three new species of *Eupenicillium*. *Trans. mycol. Soc. Japan* 9: 49-56.
- Udagawa, S., and Y. Horie. 1973a. Surface ornamentation of ascospores in *Eupenicillium* species. *Antonie van Leeuwenhoek* 39: 313-319.
- Udagawa, S., and Y. Horie. 1973b. Some *Eupenicillium* from soil of New Guinea. *Trans. mycol. Soc. Japan* 14: 370-387.
- Ueda, S. 1980. A mycofloral study on brackish water sediments in Nagasaki, Japan. *Trans. mycol. Soc. Japan* 21: 103-112. (Japanese)